

Association Between Consumption of Ultraprocessed Foods and Cognitive Decline

Natalia Gomes Gonçalves, PhD; Naomi Vidal Ferreira, PhD; Neha Khandpur, ScD; Euridice Martinez Steele, PhD; Renata Bertazzi Levy, PhD; Paulo Andrade Lotufo, MD, PhD; Isabela M. Bensenor, PhD; Paulo Caramelli, MD, PhD; Sheila Maria Alvim de Matos, PhD; Dirce M. Marchioni, PhD; Claudia Kimie Suemoto, MD, PhD

 Supplemental content

IMPORTANCE Although consumption of ultraprocessed food has been linked to higher risk of cardiovascular disease, metabolic syndrome, and obesity, little is known about the association of consumption of ultraprocessed foods with cognitive decline.

OBJECTIVE To investigate the association between ultraprocessed food consumption and cognitive decline in the Brazilian Longitudinal Study of Adult Health.

DESIGN, SETTING, AND PARTICIPANTS This was a multicenter, prospective cohort study with 3 waves, approximately 4 years apart, from 2008 to 2017. Data were analyzed from December 2021 to May 2022. Participants were public servants aged 35 to 74 years old recruited in 6 Brazilian cities. Participants who, at baseline, had incomplete food frequency questionnaire, cognitive, or covariate data were excluded. Participants who reported extreme calorie intake (<600 kcal/day or >6000 kcal/day) and those taking medication that could negatively interfere with cognitive performance were also excluded.

EXPOSURES Daily ultraprocessed food consumption as a percentage of total energy divided into quartiles.

MAIN OUTCOMES AND MEASURES Changes in cognitive performance over time evaluated by the immediate and delayed word recall, word recognition, phonemic and semantic verbal fluency tests, and Trail-Making Test B version.

RESULTS A total of 15 105 individuals were recruited and 4330 were excluded, leaving 10 775 participants whose data were analyzed. The mean (SD) age at the baseline was 51.6 (8.9) years, 5880 participants (54.6%) were women, 5723 (53.1%) were White, and 6106 (56.6%) had at least a college degree. During a median (range) follow-up of 8 (6-10) years, individuals with ultraprocessed food consumption above the first quartile showed a 28% faster rate of global cognitive decline ($\beta = -0.004$; 95% CI, -0.006 to -0.001 ; $P = .003$) and a 25% faster rate of executive function decline ($\beta = -0.003$, 95% CI, -0.005 to 0.000 ; $P = .01$) compared with those in the first quartile.

CONCLUSIONS AND RELEVANCE A higher percentage of daily energy consumption of ultraprocessed foods was associated with cognitive decline among adults from an ethnically diverse sample. These findings support current public health recommendations on limiting ultraprocessed food consumption because of their potential harm to cognitive function.

Author Affiliations: Author affiliations are listed at the end of this article.

Corresponding Author: Natalia Gomes Gonçalves, PhD, Department of Pathology, University of São Paulo Medical School, Avenida Doutor Arnaldo, 455, sala 1144, Cerqueira César, São Paulo, SP 01246-903, Brazil (natalia.g@fm.usp.br).

JAMA Neurol. doi:10.1001/jamaneurol.2022.4397
Published online December 5, 2022.

The prevalence of dementia is estimated to increase from 57 million cases in 2019 to 153 million in 2050 owing to the increase in life expectancy worldwide.^{1,2} Dementia is the most important cause of disability in high-income countries, and it is among the 10 most important causes in low-income and middle-income countries.³ The limited efficacy of available treatments for dementia highlights the importance of identifying interventions that are capable of preventing or delaying dementia onset to decrease the burden caused by this disorder.^{4,5} Lifestyle modifications, such as physical activity, healthy dietary habits, and smoking cessation, have been related to dementia prevention.⁶ Healthy eating habits, which include a high intake of whole grains, vegetables, fruit, nuts, and fish,⁷⁻¹¹ have been linked to increased brain volume and decreased risk of cognitive decline over time.^{8,9,11,12}

In the last 40 years, the food supply industries have increased the commercialization of ultraprocessed foods (UPFs).¹³ Such UPFs are formulations of processed food substances (oils, fats, sugars, starch, and protein isolates) that contain little or no whole foods and typically include flavorings, colorings, emulsifiers, and other cosmetic additives.¹⁴ Examples of UPFs are sweet and savory snacks, confectionery, breakfast cereals, ice cream, sugar-sweetened beverages, processed meats, and ready-to-eat frozen meals. Fifty-eight percent of the calories consumed by US citizens, 57% of the calories consumed by British citizens, and 48% of the calories consumed by Canadian citizens come from UPFs.¹⁵⁻¹⁷ In Brazil, this group of foods contributes to 30% of total calorie intake.¹⁸ Consumption of UPFs has been linked to an increased risk of cardiovascular disease, metabolic syndrome, and obesity.^{19,20} However, few studies have investigated the association between UPF and cognitive decline in samples from high-income countries.²¹⁻²³ Therefore, we aimed to prospectively investigate the association between UPF consumption at baseline and cognitive decline in the Brazilian Longitudinal Study of Adult Health (ELSA-Brasil).

Methods

Participants

The ELSA-Brasil is a multicenter cohort study that comprises public servants aged 35 to 74 years at baseline from 6 Brazilian cities (Belo Horizonte, Porto Alegre, Rio de Janeiro, Salvador, São Paulo, and Vitória). Data were collected in 3 waves, approximately 4 years apart, starting in 2008 to 2010. The second wave took place in 2012 to 2014, and the third wave was in 2017 to 2019. Inclusion criteria for the ELSA-Brasil study were active or retired employees of the participating institutions. Exclusion criteria were pregnancy, intention to quit working at the institution, cognitive or communication impairment, or, for those retired, residences outside of the study center area. A detailed description of the ELSA-Brasil cohort can be found elsewhere.^{24,25}

The current study excluded participants who, at baseline, did not have dietary data, had extreme amounts of energy intake (<600 kcal/day or >6000 kcal/day), had missing data on cognitive tests or covariates, or reported taking medi-

Key Points

Question Is the consumption of ultraprocessed foods associated with cognitive decline?

Findings In a cohort study of 10 775 individuals, higher consumption of ultraprocessed foods was associated with a higher rate of global and executive function decline after a median follow-up of 8 years.

Meaning These findings suggest that limiting consumption of ultraprocessed food could be associated with reduced cognitive decline in middle-aged and older adults.


cation that could negatively interfere with cognitive performance (Figure 1). A comparison between those who were included and excluded from this study analysis can be found in the eAppendix and eTable 1 in the Supplement.

This study was approved by the local ethics committees. The ELSA-Brasil study was conducted according to the guidelines of the Declaration of Helsinki,²⁶ the procedures were approved by the ethics committees of all study centers, and participants signed an informed consent before participation. This study followed the Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) reporting guideline.

Dietary Assessment

Food and drink consumption in the last 12 months was assessed at baseline using a validated Food Frequency Questionnaire (FFQ) with 114 items.²⁷ The frequency of consumption of each item was transformed into grams per day by multiplying the frequency of consumption by the weight of the portion in grams (further details can be found in the eAppendix in the Supplement). Foods were classified according to the extent of industrial processing using the Nova classification system.¹⁴ Nova includes 4 food groups. Group 1 includes unprocessed or minimally processed foods, such as fresh, dry, or frozen fruits or vegetables, grains, legumes, meat, fish, and milk, which have undergone minimal processing like grinding, roasting, pasteurization, or freezing. Group 2 includes processed culinary ingredients, such as table sugar, oils, salt, and other substances that have been extracted, pressed, or centrifuged from group 1 foods or from nature, and are used to make culinary preparations. Group 3 includes processed foods, which are manufactured using unprocessed or minimally processed foods, and ingredients from group 2 are used to prolong the durability of foods and modify their palatability. Examples of foods in group 3 include canned fruits, artisanal bread and cheese, and salted, smoked, or cured meat or fish. Group 4 includes UPFs, which are formulations of several ingredients from group 2 with food additives not used in home preparations, such as flavors, colors, sweeteners, emulsifiers, and other substances used to disguise undesirable qualities of the final product or imitate the sensorial qualities of culinary preparations from group 1.¹⁴ In the current study, foods were classified in 3 groups (eTable 2 in the Supplement): (1) unprocessed or minimally processed foods and processed culinary ingredients (Nova groups 1 and 2), (2) processed foods (Nova group 3), and

Figure 1. Flowchart of the Study Sample



ELSA indicates Brazilian Longitudinal Study of Adult Health.

(3) UPFs (Nova group 4). We calculated the daily energy consumption of UPF by summing the energy consumption in calories of all the foods in that group. Consumption of UPF was then expressed as a percentage of total daily energy consumption because a relative measure can capture the degree to which UPF composes a participant's diet while accounting for individual differences in caloric intake.

Cognitive Assessment

In this longitudinal study, individuals were tested up to 3 times every 4 years (mean [SD] time between visits, 3.3 [0.5] years). The memory domain included the immediate recall, late recall, and recognition word list tests from the Consortium to Establish a Registry for Alzheimer Disease.^{28,29} The executive function domain included the semantic and the phonemic verbal fluency tests,³⁰ and the Trail-Making Test B version.³¹ We calculated *z* scores standardized to wave 1 to compare results from different cognitive tests. A detailed description of each test and the *z* score calculation can be found in the eAppendix in the Supplement.

Covariates

Covariates that might confound the association between UPF consumption and cognitive decline included sociodemographic, clinical, and lifestyle variables. The sociodemographic variables were age, sex, monthly income per capita in US dollars, self-reported race and ethnicity (with categories self-

reported by the participants as Black or mixed [ie, mixed Black and White], White, and other races, which include Asian and Indigenous), and education (less than college and college degree or more). Race and ethnicity were assessed in this study because they are important social determinants of health that could influence cognitive performance and UPF consumption. Clinical variables included body mass index (calculated as weight in kilograms divided by height in meters squared) categories (underweight, normal weight, overweight, and obese), diabetes, hypertension, cardiovascular disease, and depression. Lifestyle factors included physical activity (light, moderate, or vigorous), smoking (never, former, or current smoker), alcohol consumption (never, former, or current alcohol use), total energy intake (in kilocalories), and adherence to a healthy diet. Details about the covariates can be found in the eAppendix in the Supplement.

Statistical Analysis

Data were analyzed from December 2021 to May 2022. Descriptive analyses were presented as mean (SD) for continuous variables and as percentages for categorical variables. We grouped the daily energy percentage contribution of UPF in quartiles (0%-19.9%, 20.0%-26.7%, 26.8%-34.1%, and 34.2%-72.7%). Our initial analysis compared each of the quartiles with the first quartile (reference group). Subsequent analysis grouped the 3 highest quartiles and compared them with the first quartile.³² We used linear mixed-effects models with random intercepts and slopes to assess the association between quartiles of UPF consumption at baseline and change in cognition over time. The timescale was the participant's age in each wave. The longitudinal association between UPF consumption and cognitive decline was evaluated by the interaction of the UPF and the timescale. The linear mixed models were adjusted for sociodemographic, clinical, and lifestyle variables. To calculate the percentage of cognitive decline rate, we subtracted each quartile slope from the first quartile slope, divided this difference by the slope of the first quartile, and multiplied it by 100. We also investigated the modifying effect of age and healthy diet scores on the association between the percentage of daily energy from UPF and cognitive decline by adding a 3-way interaction among the percentage of daily energy from UPF, the timescale, and each modifier on our main models. The significant interactions were assessed in stratified analyses. Inverse probability weighting (IPW) was used to account for nonresponse across waves.^{33,34} Details about the IPW calculation can be found in the Supplement.

We performed 2 sets of sensitivity analyses. First, we repeated the analysis by excluding participants who reported caloric intakes above the 95th percentile (5831 kcal for men and 4607 kcal for women) and without excluding participants because of caloric intake. Moreover, we verified the robustness of our findings despite the missing cognitive data by design in wave 2 by imputing cognitive data using next observation carried backward using the scores from wave 3 in wave 2 for those who were younger than 55 years in wave 2. We assumed this approach is conservative since cognitive performance is expected to decline over time.³⁵ The α level was set at the 5% level

in 2-sided tests. Statistical analyses were performed using R statistical software version 3.6.3 (R Project for Statistical Computing) using the lme4 package.^{36,37}

Results

Sample Characteristics

A total of 15 105 individuals were recruited and 4330 were excluded, leaving 10 775 participants. The median (range) duration of follow-up was 8 (6-10) years. At baseline, the mean (SD) age of the participants was 51.6 (8.9) years, 5880 participants (54.6%) were women, 5723 (53.1%) were

White, and 6106 (56.6%) had at least a college education. The mean (SD) BMI was 26.9 (4.7), and the mean (SD) total daily calorie intake was 2856 (992) kcal, 27% of which came from UPF (mean [SD], 785.0 [419.1] kcal/day). Compared with the lower quartile of the percentage of daily energy from UPF, those in the fourth quartile (ie, highest UPF consumption) were more likely to be younger, women, White, had higher education and income, were more likely to be nonsmokers, and less likely to be current alcohol consumers. The highest quartile also had a higher total energy intake, lower physical activity, and lower frequency of comorbidities, but a higher frequency of depressive symptoms (Table 1).

Table 1. Baseline Characteristics of the Study Sample by Quartiles of the Percentage of Daily Energy From UPFs

Characteristic	Participants, No. (%)					P value
	Total (N = 10 775)	UPF consumption quartile ^a				
		1 (0%-19.9%) (n = 2694)	2 (20.0%-26.7%) (n = 2694)	3 (26.8%-34.1%) (n = 2694)	4 (34.2%-72.7%) (n = 2693)	
Age, mean (SD), y	51.6 (8.9)	54.2 (8.6)	52.0 (9.0)	50.8 (9.0)	49.6 (9.0)	<.001
Sex						
Female	5880 (54.6)	1392 (51.6)	1443 (53.5)	1501 (55.7)	1544 (57.3)	<.001
Male	4895 (45.4)	1302 (48.4)	1251 (46.5)	1193 (44.3)	1149 (42.7)	
Self-reported race						
Black or mixed ^b	4685 (43.5)	1404 (52.1)	1215 (45.3)	1069 (39.7)	1047 (36.9)	<.001
White	5723 (53.1)	1168 (43.3)	1384 (51.4)	1542 (57.2)	1542 (60.5)	
Other ^c	367 (3.4)	122 (4.6)	95 (3.8)	83 (3.1)	67 (2.6)	
Education (less than college degree)	4669 (43.4)	1390 (51.6)	1154 (42.9)	1037 (38.5)	1088 (40.4)	<.001
Monthly income, mean (SD), \$US	983.6 (788.1)	953.5 (797.8)	974.3 (785.8)	997.8 (776.5)	1010.4 (791.8)	.004
Body mass index, mean (SD) ^d	26.9 (4.7)	26.9 (4.6)	27.0 (4.6)	27.0 (4.8)	27.0 (4.8)	.25
Total calorie intake, mean (SD), kcal	2855.9 (991.7)	2813.0 (977.0)	2858.0 (999.0)	2860.0 (993.0)	2894.0 (997.0)	.004
Calories from ultraprocessed foods, mean (SD), kcal	785.0 (419.1)	416.0 (185.0)	667.0 (241.0)	865.0 (308.0)	1192.0 (440.0)	<.001
Calories from Nova groups, % of total calories ^e						
Groups 1 and 2	1873.5 (65.6)	2194.1 (78.0)	1986.3 (69.5)	1790.3 (62.6)	1522.2 (52.6)	<.001
Group 3	197.4 (7.0)	202.9 (7.4)	208.7 (7.3)	203.2 (7.3)	179.8 (6.2)	
Group 4	785.0 (27.4)	416.0 (14.8)	663.0 (23.3)	866.5 (30.3)	1192.0 (41.2)	
Physical activity						
None or light	8197 (76.1)	1968 (73.0)	1996 (74.1)	2062 (76.5)	2171 (80.6)	<.001
Moderate	1539 (14.3)	458 (17.0)	406 (15.1)	383 (14.2)	292 (11.0)	
Vigorous	1039 (9.6)	268 (10.0)	292 (10.8)	249 (9.3)	230 (8.4)	
Hypertension, yes	3704 (34.4)	1100 (41.0)	969 (36.0)	813 (30.2)	822 (30.5)	<.001
Diabetes, yes	2016 (18.7)	701 (26.0)	508 (19.0)	424 (15.7)	383 (14.2)	<.001
Cardiovascular disease, yes	578 (5.3)	165 (6.1)	153 (5.7)	139 (5.1)	121 (4.5)	.04
Depressive symptoms, yes	1263 (11.7)	295 (11.0)	306 (11.3)	285 (10.6)	377 (14.0)	<.001
Alcohol consumption						
Never	1061 (9.8)	301 (11.2)	248 (9.2)	259 (9.6)	253 (9.3)	<.001
Former	1967 (18.3)	467 (17.3)	439 (16.3)	475 (17.7)	586 (21.7)	
Current	7747 (71.9)	1926 (71.5)	2007 (74.5)	1960 (72.7)	1854 (69.0)	
Smoking						
Never	6297 (58.4)	1474 (54.7)	1527 (56.7)	1633 (60.6)	1663 (61.7)	<.001
Former	3165 (29.4)	847 (31.4)	851 (31.6)	748 (27.7)	719 (26.7)	
Current	1313 (12.2)	373 (13.9)	316 (11.7)	313 (11.7)	311 (11.6)	

Abbreviation: UPF, ultraprocessed food.

^a UPFs are represented as a percentage of total daily energy consumption divided in quartiles.

^b Includes mixed Black and White.

^c Includes Asian, Indigenous, or other ethnic groups.

^d Body mass index calculated as weight in kilograms divided by height in meters squared.

^e Group 1 includes unprocessed or minimally processed foods. Group 2 includes processed culinary ingredients. Group 3 includes processed foods. Group 4 includes UPFs.

Table 2. Association Between Baseline Quartiles of the Percentage of Daily Energy From Ultraprocessed Foods and Yearly Cognitive Change During the Study Period

Domain	Change in standardized cognitive score per year						Difference, % ^d
	Model 1 ^a		Model 2 ^b		Model 3 ^c		
	β (95% CI)	P value for trend	β (95% CI)	P value for trend	β (95% CI)	P value for trend	
Memory							
All quartiles							
Quartile 1 × time	0 [Reference]		0 [Reference]		0 [Reference]		0 [Reference]
Quartile 2 × time	0.001 (−0.003 to 0.004)		0.001 (−0.004 to 0.006)		0.001 (−0.002 to 0.004)		6
Quartile 3 × time	0.000 (−0.004 to 0.003)	.88	0.000 (−0.005 to 0.004)	.82	0.000 (−0.004 to 0.003)	.80	0
Quartile 4 × time	0.001 (−0.002 to 0.004)		0.002 (−0.003 to 0.006)		0.001 (−0.002 to 0.005)		6
Lowest × highest quartiles							
Quartile 1 (lowest 25%)	0 [Reference]		0 [Reference]		0 [Reference]		0 [Reference]
Quartiles 2-4 (highest 75%)	0.000 (−0.003 to 0.003)	.86	0.000 (−0.002 to 0.003)	.77	0.000 (−0.002 to 0.003)	.77	0
Executive function							
All quartiles							
Quartile 1 × time	0 [Reference]		0 [Reference]		0 [Reference]		0 [Reference]
Quartile 2 × time	−0.003 (−0.006 to 0.000)		−0.003 (−0.006 to 0.000)		−0.003 (−0.006 to 0.000)		25
Quartile 3 × time	−0.002 (−0.005 to 0.001)	.23	−0.003 (−0.006 to 0.000)	.12	−0.003 (−0.005 to 0.000)	.12	25
Quartile 4 × time	−0.002 (−0.005 to 0.001)		−0.002 (−0.005 to 0.001)		−0.002 (−0.005 to 0.001)		16
Lowest × highest quartiles							
Quartile 1 (lowest 25%)	0 [Reference]		0 [Reference]		0 [Reference]		0 [Reference]
Quartiles 2-4 (highest 75%)	−0.002 (−0.005 to 0.000)	.04	−0.003 (−0.005 to 0.000)	.01	−0.003 (−0.005 to 0.000)	.01	25
Global cognition							
All quartiles							
Quartile 1 × time	0 [Reference]		0 [Reference]		0 [Reference]		0 [Reference]
Quartile 2 × time	−0.003 (−0.006 to 0.000)		−0.003 (−0.006 to 0.000)		−0.003 (−0.006 to 0.000)		21
Quartile 3 × time	−0.004 (−0.007 to −0.001)	.06	−0.004 (−0.007 to −0.001)	.04	−0.004 (−0.007 to −0.001)	.04	28
Quartile 4 × time	−0.003 (−0.006 to 0.000)		−0.003 (−0.006 to 0.000)		−0.003 (−0.006 to 0.000)		21
Lowest × highest quartiles							
Quartile 1 (lowest 25%)	0 [Reference]		0 [Reference]		0 [Reference]		0 [Reference]
Quartiles 2-4 (highest 75%)	−0.003 (−0.005 to −0.001)	.004	−0.003 (−0.006 to −0.001)	.003	−0.003 (−0.006 to −0.001)	.004	28

^a Model 1 includes linear mixed models adjusted for age, sex, race and ethnicity, education, and income.

^b Model 2 includes linear mixed models additionally adjusted for physical activity, body mass index, hypertension, diabetes, cardiovascular disease, depressive symptoms, alcohol consumption, and smoking.

^c Model 3 includes linear mixed models additionally adjusted for total calories and healthy eating score.


^d Difference is between each quartile and the first quartile in model 3.

Cognitive Performance and Consumption of UPFs

After a median follow-up of 8 years, participants who reported consumption of UPF of more than 19.9% of daily calories had a 28% faster rate of global cognitive decline compared with those who reported consumption of UPF up to 19.9% of daily calories (β = −0.004; 95% CI, −0.006 to −0.001; P = .003) (Table 2 and Figure 2). Moreover, participants who reported consumption of UPF more than 19.9% of daily calo-

ries had a 25% faster rate of executive function decline compared with those who reported consumption of UPF less than or equal to 19.9% of daily calories (β = −0.003, 95% CI, −0.005 to 0.000; P = .01) (Table 2). We found no association between the percentage of daily energy from UPF and the memory score. Age was an effect modifier in the association of the percentage of daily energy from UPF and cognitive function (P for interaction < .001). Participants younger than 60

Figure 2. Trajectories of Global Cognitive Performance Over Time According to Quartiles of the Percentage of Daily Energy From Ultraprocessed Foods (%UPF)



Mixed linear regression models with random intercepts and slopes were adjusted for age, sex, race and ethnicity, education, income, physical activity, body mass index, hypertension, diabetes, cardiovascular disease, depressive symptoms, alcohol consumption, smoking, total calories, and healthy eating score. *P* values were calculated for the interaction between UPF quartiles (ordinal continuous variable) and age as the timescale. Shaded areas indicate 95% CIs.

years with UPF consumption greater than 19.9% showed a faster global cognition decline compared with those with UPF consumption less than 19.9% ($\beta = -0.006$; 95% CI, -0.009 to -0.003 ; $P < .001$) (eTable 3 in the Supplement), whereas there was no association of the percentage of daily energy from UPF and global cognition decline for those aged 60 years or older (eTable 3 in the Supplement). Adherence to a healthy diet was also an effect modifier on the association of UPF and global cognitive function (*P* for interaction = .04). Participants with low healthy diet scores who consumed more than 19.9% of calories from UPF showed a faster global cognition decline compared with those who consumed less than 19.9% ($\beta = -0.005$; 95% CI, -0.009 to -0.002 ; $P = .004$) (eTable 4 in the Supplement and Figure 3A). We found no association between the percentage of daily energy from UPF and global cognition for participants with high healthy diet scores (eTable 4 in the Supplement and Figure 3B).

To assess the robustness of our findings, we performed sensitivity analyses excluding participants who reported caloric intakes above the 95th percentile by sex and without excluding participants because of caloric intake. We also imputed data for cognitive performance for participants younger than 55 years in wave 2. The results of the sensitivity analyses were similar to those found in the main analysis (eTable 5 and eTable 6 in the Supplement).

Discussion

In this cohort study of 10 775 individuals followed for a median of 8 years, we found that consumption of UPF greater than 19.9% of total daily calories was associated with a faster decline in global cognitive performance and executive function

compared with consumption less than 19.9% of total daily calories. We also found that the percentage of daily energy from UPF was associated with cognitive decline in participants younger than 60 years, which suggests the importance of preventive interventions in middle-aged adults. Additionally, the percentage of daily energy from UPF was associated with cognitive decline in participants with a low healthy diet score, whereas there was no association in those with a high healthy diet score. Our findings are in line with previous studies linking consumption of UPF and adverse health outcomes, such as the increased risk of overweight and obesity,^{38,39} metabolic syndrome,⁴⁰ cancer,⁴¹ cardiovascular diseases,⁴² and all-cause mortality.^{43,44}


A prior study²¹ that investigated the association of UPF and cognition in 568 individuals with type 2 diabetes found no association between total calorie consumption of UPF and cognitive decline, likely because of the small sample size or reverse causation, since individuals who develop diabetes could have reduced their UPF consumption after diagnosis. Our findings are in line with 2 recent studies^{22,23} that investigated the association between UPF consumption and cognition. A cross-sectional study²² of older US adults found an association between UPF consumption and worse verbal fluency performance in participants without preexisting chronic health conditions. Another study²³ investigated the association between consumption of UPF and dementia in 72 083 adults aged 55 years or older from the UK Biobank. Consumption of UPF was associated with a higher risk of all-cause dementia, vascular dementia, and Alzheimer disease after 10 years of follow-up.²³ The association between UPF and cognitive decline found in our study, particularly the decline in executive function, could be secondary to cerebrovascular lesions resulting from UPF consumption, because these functions are particularly sensitive to microvascular lesions.^{42,45}

Neuroimaging studies^{46,47} have found that high consumption of a Western dietary pattern was related to a reduction in the left hippocampus and gray matter volume in cognitively healthy individuals. Another possible biological mechanism for the decline in executive function and global cognition seen in our study may be related to systemic inflammation caused by the consumption of UPF, because increased levels of circulating proinflammatory cytokines have been associated with cognitive decline.⁴⁸⁻⁵² On the other hand, healthy dietary patterns were associated with higher gray and white matter volume, total brain volume, and A β 42/40 ratio, as well as lower oxidative stress and inflammation,^{53,54} which could explain our findings that the percentage of daily energy from UPF was associated with cognitive decline in participants with a low healthy diet score, but not in those with a high healthy diet score.

Strengths and Limitations

This study has some strengths. First, this is a large ethnically diverse cohort study from a low- and middle-income country with up to 10 years of follow-up. Second, the diet assessment was conducted using a validated questionnaire.²⁷ Moreover, we found that UPF consumption was associated with cognitive decline in middle-aged participants. The inclusion of

Figure 3. Trajectories of Global Cognitive Performances Over Time in Participants With Low Healthy Diet Scores and High Healthy Diet Scores



Graphs show trajectories of global cognitive performances over time according to quartiles of the percentage of daily energy from ultraprocessed foods (%UPF). Mixed linear regression models with random intercepts and slopes were adjusted for age, sex, race and ethnicity, education, income, physical activity, body mass index, hypertension, diabetes, cardiovascular disease, depressive symptoms, alcohol consumption, smoking, total calories, and healthy eating score. Shaded areas indicate 95% CIs.

middle-aged participants in studies about risk factors for cognitive decline is particularly important to understand potential preventive targets early in the life course.^{55,56}

However, our findings should be considered in light of study limitations. Attrition is a concern for a long-term study, and participants younger than 55 years were not submitted to cognitive assessment during the second visit, because of the study design. Nevertheless, the use of IPW provided some correction for selection bias. Moreover, the imputation of missing cognitive data by design in wave 2 did not change the main study findings. In addition, diet was assessed only at baseline, which may not reflect longitudinal diet changes and may lead to an underestimation of the associations between UPF and cognition.⁵⁷ A few food items may have been misclassified because the FFQ was not specifically designed to assess the degree of processing. Because the FFQ was self-reported, the UPF consumption could be underreported owing to social desirability bias, which could have biased associations toward the null. Using the same calorie cutoff interval for men and women may introduce bias due to different caloric intake needs. However, the use of a relative measure as the exposure variable (percentage of the daily energy from UPF) minimizes the effect of extreme total energy intakes on the studied association.⁵⁸ Additionally, a sensitivity analysis using different cutoffs for calorie intake in men and women showed similar results to our main analysis. Although the use of software based on North American foods to estimate calorie content may be a limitation, it is unlikely to bias the UPF con-

sumption estimations, because the UPF classification used the composition of products commonly consumed in Brazil and did not consider the nutritional composition from the software. Additionally, our findings may be subjected to selection bias, because the characteristics of those included and those excluded in the study at baseline differed. Although we adjusted the analyses for several sociodemographic and clinical confounders, we cannot exclude the possibility of residual confounding. Furthermore, since neuroimaging is not available in the ELSA-Brasil study, we were not able to investigate possible mechanisms that could explain the association between UPF consumption and cognitive decline in our study.

Conclusions

In this large cohort study, a higher percentage of daily energy from UPF was associated with cognitive decline during 8 years of follow-up. Intact cognitive function is key to successful aging. Therefore, despite the small effect size of the association between UPF consumption and cognitive decline, our findings are meaningful to cognitive health. Limiting UPF consumption, particularly in middle-aged adults, may be an efficient form to prevent cognitive decline. Future studies investigating the mechanism by which UPF may lead to cognitive decline are needed, as well as confirmation of our findings in other longitudinal studies and randomized clinical trials.

ARTICLE INFORMATION

Accepted for Publication: October 7, 2022.

Published Online: December 5, 2022.
doi:10.1001/jamaneurol.2022.4397

Author Affiliations: Department of Pathology, University of São Paulo Medical School, São Paulo, Brazil (Gomes Gonçalves); Adventist University of

São Paulo, Engenheiro Coelho, Brazil (Vidal Ferreira); Division of Geriatrics, University of São Paulo Medical School, São Paulo, Brazil (Vidal Ferreira, Suemoto); Department of Nutrition, School of Public Health, University of São Paulo, São Paulo, Brazil (Khandpur, Martinez Steele, Marchioni); Department of Nutrition, Harvard T.H. Chan School of Public Health, Boston,

Massachusetts (Khandpur); Department of Preventive Medicine, School of Medicine, University of São Paulo, São Paulo, Brazil (Bertazzi Levy); Center for Clinical and Epidemiological Research, Hospital Universitário, University of São Paulo, São Paulo, Brazil (Andrade Lotufo, Bensenor); Behavioral and Cognitive Neurology Research Unit, Faculdade de Medicina,

Universidade Federal de Minas Gerais, Belo Horizonte, Brazil (Caramelli); Federal University of Bahia, Salvador, Brazil (Alvim de Matos).

Author Contributions: Dr Gomes Gonçalves had full access to all of the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis.

Concept and design: Gomes Gonçalves, Khandpur, Andrade Lotufo, Caramelli, Alvim de Matos, Suemoto.

Acquisition, analysis, or interpretation of data: Gomes Gonçalves, Vidal Ferreira, Khandpur, Martinez Steele, Bertazzi Levy, Andrade Lotufo, Bensenor, Alvim de Matos, Marchioni, Suemoto. **Drafting of the manuscript:** Gomes Gonçalves, Martinez Steele, Suemoto.

Critical revision of the manuscript for important intellectual content: All authors.

Statistical analysis: Gomes Gonçalves, Vidal Ferreira, Khandpur, Martinez Steele, Bertazzi Levy. **Obtained funding:** Andrade Lotufo, Bensenor, Alvim de Matos.

Administrative, technical, or material support: Khandpur, Martinez Steele, Bertazzi Levy, Andrade Lotufo, Bensenor, Alvim de Matos, Suemoto. **Supervision:** Andrade Lotufo, Bensenor, Alvim de Matos, Suemoto.

Conflict of Interest Disclosures: None reported.

Funding/Support: The ELSA-Brasil study was supported by the Brazilian Ministry of Health, the Ministry of Science, Technology and Innovation and the National Council for Scientific and Technological Development (CNPq). Drs Bertazzi Levy, Andrade Lotufo, Bensenor, Caramelli, Marchioni, and Suemoto receive support from CNPq, Brazil (research productivity fellowship).

Role of the Funder/Sponsor: The funders had no role in the design and conduct of the study; collection, management, analysis, and interpretation of the data; preparation, review, or approval of the manuscript; and decision to submit the manuscript for publication.

REFERENCES

- Nichols E, Steinmetz JD, Vollset SE, et al; GBD 2019 Dementia Forecasting Collaborators. Estimation of the global prevalence of dementia in 2019 and forecasted prevalence in 2050: an analysis for the Global Burden of Disease Study 2019. *Lancet Public Health*. 2022;7(2):e105-e125. doi:10.1016/S2468-2667(21)00249-8
- World Health Organization. Risk reduction of cognitive decline and dementia: WHO guidelines. Accessed October 28, 2022. <https://apps.who.int/iris/bitstream/handle/10665/312180/9789241550543-eng.pdf>
- World Health Organization. *World Report on Ageing and Health 2015*. World Health Organization; 2015.
- Feldman HH, Haas M, Gandy S, et al; One Mind for Research and the New York Academy of Sciences. Alzheimer's disease research and development: a call for a new research roadmap. *Ann N Y Acad Sci*. 2014;1313(1):1-16. doi:10.1111/nyas.12424
- Barnes DE, Yaffe K. The projected effect of risk factor reduction on Alzheimer's disease prevalence. *Lancet Neurol*. 2011;10(9):819-828. doi:10.1016/S1474-4422(11)70072-2
- Prince M, Albanese E, Guerchet M, Prina M; Alzheimer's Disease International. Dementia and risk reduction: an analysis of protective and modifiable factors. September 2014. Accessed October 28, 2022. <https://www.alzint.org/u/WorldAlzheimerReport2014.pdf>
- Ekstrand B, Scheers N, Rasmussen MK, Young JF, Ross AB, Landberg R. Brain foods: the role of diet in brain performance and health. *Nutr Rev*. 2021;79(6):693-708. doi:10.1093/nutrit/nuaa091
- Croll PH, Voortman T, Ikram MA, et al. Better diet quality relates to larger brain tissue volumes: the Rotterdam Study. *Neurology*. 2018;90(24):e2166-e2173. doi:10.1212/WNL.0000000000005691
- Morris MC, Tangney CC, Wang Y, et al. MIND diet slows cognitive decline with aging. *Alzheimers Dement*. 2015;11(9):1015-1022. doi:10.1016/j.jalz.2015.04.011
- Scarmeas N, Stern Y, Tang MX, Mayeux R, Luchsinger JA. Mediterranean diet and risk for Alzheimer's disease. *Ann Neurol*. 2006;59(6):912-921. doi:10.1002/ana.20854
- Morris MC, Evans DA, Tangney CC, Bienias JL, Wilson RS. Fish consumption and cognitive decline with age in a large community study. *Arch Neurol*. 2005;62(12):1849-1853. doi:10.1001/archneur.62.12.noc50161
- Scarmeas N, Stern Y, Mayeux R, Manly JJ, Schupf N, Luchsinger JA. Mediterranean diet and mild cognitive impairment. *Arch Neurol*. 2009;66(2):216-225. doi:10.1001/archneurol.2008.536
- Popkin BM, Adair LS, Ng SW. Global nutrition transition and the pandemic of obesity in developing countries. *Nutr Rev*. 2012;70(1):3-21. doi:10.1111/j.1753-4887.2011.00456.x
- Monteiro CA, Cannon G, Levy RB, et al. Ultra-processed foods: what they are and how to identify them. *Public Health Nutr*. 2019;22(5):936-941. doi:10.1017/S1368980018003762
- Martinez Steele E, Baraldi LG, Louzada ML, Moubarac JC, Mozaffarian D, Monteiro CA. Ultra-processed foods and added sugars in the US diet: evidence from a nationally representative cross-sectional study. *BMJ Open*. 2016;6(3):e009892. doi:10.1136/bmjopen-2015-009892
- Rauber F, Louzada MLDC, Martinez Steele E, et al. Ultra-processed foods and excessive free sugar intake in the UK: a nationally representative cross-sectional study. *BMJ Open*. 2019;9(10):e027546. doi:10.1136/bmjopen-2018-027546
- Moubarac JC, Batal M, Louzada ML, Martinez Steele E, Monteiro CA. Consumption of ultra-processed foods predicts diet quality in Canada. *Appetite*. 2017;108:512-520. doi:10.1016/j.appet.2016.11.006
- Louzada ML da C, Baraldi LG, Steele EM, et al. Consumption of ultra-processed foods and obesity in Brazilian adolescents and adults. *Prev Med*. 2015; 81:9-15. doi:10.1016/j.jypmed.2015.07.018
- Pagliai G, Dinu M, Madarena MP, Bonaccio M, Iacoviello L, Sofi F. Consumption of ultra-processed foods and health status: a systematic review and meta-analysis. *Br J Nutr*. 2021;125(3):308-318. doi:10.1017/S0007114520002688
- Costa de Miranda R, Rauber F, Levy RB. Impact of ultra-processed food consumption on metabolic health. *Curr Opin Lipidol*. 2021;32(1):24-37. doi:10.1097/MOL.0000000000000728
- Weinstein G, Vered S, Ivancovsky-Wajcman D, et al. Consumption of ultra-processed food and cognitive decline among older adults with type-2 diabetes. *J Gerontol Ser A*. Published online March 19, 2022. doi:10.1093/gerona/glac070
- Cardoso BR, Machado P, Martinez Steele E. Association between ultra-processed food consumption and cognitive performance in US older adults: a cross-sectional analysis of the NHANES 2011-2014. *Eur J Nutr*. Published online July 1, 2022. doi:10.1007/s00394-022-02911-1
- Li H, Li S, Yang H, et al. Association of ultraprocessed food consumption with risk of dementia: a prospective cohort study. *Neurology*. 2022;99(10):e1056-e1066. doi:10.1212/WNL.0000000000200871
- Aquino EML, Barreto SM, Bensenor IM, et al. Brazilian Longitudinal Study of Adult Health (ELSA-Brasil): objectives and design. *Am J Epidemiol*. 2012;175(4):315-324. doi:10.1093/aje/kwr294
- Schmidt MI, Duncan BB, Mill JG, et al. Cohort profile: Longitudinal Study of Adult Health (ELSA-Brasil). *Int J Epidemiol*. 2015;44(1):68-75. doi:10.1093/ije/dyu027
- World Medical Association. World Medical Association Declaration of Helsinki: ethical principles for medical research involving human subjects. *JAMA*. 2013;310(20):2191-2194. doi:10.1001/jama.2013.281053
- Molina MDCB, Benseñor IM, de Oliveria Cardoso L, et al. Reproducibility and relative validity of the Food Frequency Questionnaire used in the ELSA-Brasil [in Portuguese]. *Cad Saude Publica*. 2013;29:2. doi:10.1590/S0102-311X2013000600024
- Morris JC, Heyman A, Mohs RC, et al. The Consortium to Establish a Registry for Alzheimer's disease (CERAD). Part 1. Clinical and neuropsychological assessment of Alzheimer's disease. *Neurology*. 1989;39(9):1159-1165. doi:10.1212/WNL.39.9.1159
- Bertolucci PHF, Okamoto IH, Toniolo Neto J, Ramos LR, Brucki SMD. Desempenho da população brasileira na bateria neuropsicológica do Consortium to Establish a Registry for Alzheimer's Disease (CERAD). *Arch Clin Psychiatry*. 1998. Accessed October 28, 2022. <https://www.scienceopen.com/document?vid=74d8c494-f785-43f4-a54b-5c7b4e05219d>
- Strauss E, Sherman E, Spreen O. *A Compendium of Neuropsychological Tests: Administration, Norms, and Commentary*. 3rd Edition. Oxford University Press; 2006.
- Greenleaf CL, Margolis RB, Erker GJ. Application of the Trail Making Test in differentiating neuropsychological impairment of elderly persons. *Percept Mot Skills*. 1985;61(3 Pt 2):1283-1289. doi:10.2466/pms.1985.61.3f.1283
- Chiesa ST, Masi S, Shipley MJ, et al. Carotid artery wave intensity in mid- to late-life predicts cognitive decline: the Whitehall II study. *Eur Heart J*. 2019;40(28):2300-2309. doi:10.1093/eurheartj/ehz189
- Hernán MA, Robins JM. Estimating causal effects from epidemiological data. *J Epidemiol Community Health*. 2006;60(7):578-586. doi:10.1136/jech.2004.029496
- Weuve J, Tchetgen Tchetgen EJ, Glymour MM, et al. Accounting for bias due to selective attrition: the example of smoking and cognitive decline.

- Epidemiology*. 2012;23(1):119-128. doi:10.1097/EDE.0b013e318230e861
35. Engels JM, Diehr P. Imputation of missing longitudinal data: a comparison of methods. *J Clin Epidemiol*. 2003;56(10):968-976. doi:10.1016/S0895-4356(03)00170-7
36. R Development Core Team. R: A language and environment for statistical computing. R Foundation for Statistical Computing. 2020. Accessed October 28, 2022. <https://www.r-project.org/>
37. Bates D, Machler M, Bolker B, et al. R package "lme4." 2021. Accessed October 28, 2022. <https://cran.r-project.org/web/packages/lme4/lme4.pdf>
38. Mendonça RD, Pimenta AM, Gea A, et al. Ultra-processed food consumption and risk of overweight and obesity: the University of Navarra Follow-Up (SUN) cohort study. *Am J Clin Nutr*. 2016;104(5):1433-1440. doi:10.3945/ajcn.116.135004
39. Canhada SL, Luft VC, Giatti L, et al. Ultra-processed foods, incident overweight and obesity, and longitudinal changes in weight and waist circumference: the Brazilian Longitudinal Study of Adult Health (ELSA-Brasil). *Public Health Nutr*. 2020;23(6):1076-1086. doi:10.1017/S13688980190002854
40. Martínez Steele E, Juul F, Neri D, Rauber F, Monteiro CA. Dietary share of ultra-processed foods and metabolic syndrome in the US adult population. *Prev Med*. 2019;125:40-48. doi:10.1016/j.jpmed.2019.05.004
41. Fiolet T, Srour B, Sellem L, et al. Consumption of ultra-processed foods and cancer risk: results from NutriNet-Santé prospective cohort. *BMJ*. 2018;360:k322. doi:10.1136/bmj.k322
42. Srour B, Fezeu LK, Kesse-Guyot E, et al. Ultra-processed food intake and risk of cardiovascular disease: prospective cohort study (NutriNet-Santé). *BMJ*. 2019;365:11451. doi:10.1136/bmj.11451
43. Rico-Campà A, Martínez-González MA, Alvarez-Alvarez I, et al. Association between consumption of ultra-processed foods and all cause mortality: SUN prospective cohort study. *BMJ*. 2019;365:11949. doi:10.1136/bmj.11949
44. Kim H, Hu EA, Rebholz CM. Ultra-processed food intake and mortality in the USA: results from the Third National Health and Nutrition Examination Survey (NHANES III, 1988-1994). *Public Health Nutr*. 2019;22(10):1777-1785. doi:10.1017/S1368898018003890
45. Jokinen H, Kalska H, Mäntylä R, et al. Cognitive profile of subcortical ischaemic vascular disease. *J Neurol Neurosurg Psychiatry*. 2006;77(1):28-33. doi:10.1136/jnnp.2005.069120
46. Jacka FN, Cherbuin N, Anstey KJ, Sachdev P, Butterworth P. Western diet is associated with a smaller hippocampus: a longitudinal investigation. *BMC Med*. 2015;13(1):215. doi:10.1186/s12916-015-0461-x
47. Berti V, Murray J, Davies M, et al. Nutrient patterns and brain biomarkers of Alzheimer's disease in cognitively normal individuals. *J Nutr Health Aging*. 2015;19(4):413-423. doi:10.1007/s12603-014-0534-0
48. Shen W, Wolf PG, Carbonero F, et al. Intestinal and systemic inflammatory responses are positively associated with sulfidogenic bacteria abundance in high-fat-fed male C57BL/6J mice. *J Nutr*. 2014;144(8):1181-1187. doi:10.3945/jn.114.194332
49. Beilharz JE, Maniam J, Morris MJ. Short exposure to a diet rich in both fat and sugar or sugar alone impairs place, but not object recognition memory in rats. *Brain Behav Immun*. 2014;37:134-141. doi:10.1016/j.bbi.2013.11.016
50. Ghanim H, Abuaysheh S, Sia CL, et al. Increase in plasma endotoxin concentrations and the expression of Toll-like receptors and suppressor of cytokine signaling-3 in mononuclear cells after a high-fat, high-carbohydrate meal: implications for insulin resistance. *Diabetes Care*. 2009;32(12):2281-2287. doi:10.2337/dc09-0979
51. Schram MT, Euser SM, de Craen AJM, et al. Systemic markers of inflammation and cognitive decline in old age. *J Am Geriatr Soc*. 2007;55(5):708-716. doi:10.1111/j.1532-5415.2007.01159.x
52. Calice-Silva V, Suemoto CK, Brunoni AR, Bensenor IM, Lotufo PA. Association between GlycA and cognitive function cross-sectional results from the ELSA-Brasil study. *Alzheimer Dis Assoc Disord*. 2021;35(2):128-134. doi:10.1097/WAD.0000000000000431
53. Gu Y, Brickman AM, Stern Y, et al. Mediterranean diet and brain structure in a multiethnic elderly cohort. *Neurology*. 2015;85(20):1744-1751. doi:10.1212/WNL.0000000000002121
54. Bagetta D, Maruca A, Lupia A, et al. Mediterranean products as promising source of multi-target agents in the treatment of metabolic syndrome. *Eur J Med Chem*. 2020;186:111903. doi:10.1016/j.ejmech.2019.111903
55. Sperling RA, Aisen PS, Beckett LA, et al. Toward defining the preclinical stages of Alzheimer's disease: recommendations from the National Institute on Aging-Alzheimer's Association workgroups on diagnostic guidelines for Alzheimer's disease. *Alzheimers Dement*. 2011;7(3):280-292. doi:10.1016/j.jalz.2011.03.003
56. Livingston G, Huntley J, Sommerlad A, et al. Dementia prevention, intervention, and care: 2020 report of the Lancet Commission. *Lancet*. 2020;396(10248):413-446. doi:10.1016/S0140-6736(20)30367-6
57. Clarke R, Shipley M, Lewington S, et al. Underestimation of risk associations due to regression dilution in long-term follow-up of prospective studies. *Am J Epidemiol*. 1999;150(4):341-353. doi:10.1093/oxfordjournals.aje.a010013
58. Willett W. *Nutritional Epidemiology*. Oxford Academic Press; 2013. doi:10.1093/acprof:oso/9780199754038.001.0001